



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
REQUEST FOR FILING REISSUE PATENT APPLICATION

AIRE

The Commissioner of Patents and Trademarks
Washington, D.C. 20231



Sir:

This is a request for a reissue patent application of:

Inventor: Robert H. Reid, John E. van Hamont, William R. Brown,
Edgar C.Boedeker, Curt Thies

Title: MICROPARTICLE CARRIERS OF MAXIMAL UPTAKE CAPACITY
BY BOTH M CELLS AND NON-M CELLS

Patent No: 5,693,343 **Issue Date:** December 2, 1997

Serial No.: 08/242,960 **Filing Date:** May 16, 1994

Date of this reissue application filing: November 30, 1999

Attorney docket no: Army 103

This application for reissue includes:

X Specification: 15 pages (only spec. and claims)

X Abstract: 1 page(s)

X Drawings:* sheet(s) per set:

1 set(s) informal (Figs. 1-19);

set(s) formal of size A4 13" 14"

*see separate letter requesting transfer of the drawings from the patent file

X Declaration

X Assent of Assignee

Request for Title Search

Verified Statement establishing small entity status

X Power of Attorney (included in Declaration)

X Also attached:

(1) Form PTO-1449

(2) copies of references cited

(3) Letter Requesting Transfer of Drawings from the Patent File

09451321

11/30/99
1545 U.S. PRO

Reid, et al. Reissue Application of U.S. Patent 5,693,343

Priority is claimed under 37 CFR 1.78(a)(3) based on: Ser. No. 867,301, filed April 10, 1992, Pat. No. 5,417,986 which is a continuation in part of Ser. No. 805,721, filed Nov. 21, 1991, abandoned, which is a continuation in part of Ser. No. 690,485, filed April 24, 1991, abandoned, which is a continuation in part of Ser. No. 521,945, filed May 11, 1990, abandoned, which is a continuation in part of Ser. No. 493,597, filed March 15, 1990, abandoned, which is a continuation in part of Ser. No. 590,308, filed March 16, 1984.

Fee Calculation:

Basic Filing Fee (\$760/\$380) \$ 760

Claim fees:

	<u>Claims in</u> <u>reissue app.</u>	<u>Claims in</u> <u>orig. patent</u>	
Total effective claims	(A) 24	(B) 7	(A) minus [larger of (B) or 20] = <u>4</u> x \$18/\$9= \$72.00
Total indep. claims	(D) 7	(E) 1	(D) minus (E) = <u>6</u> x \$78/\$39= \$468.00

Title Search Fee (\$25.00) \$

Petition Fee per Rule 17(h) for Rule 47 Petition (\$130.00) \$

Other Fees: \$

TOTAL FEES \$1300.00

CHARGE STATEMENT: The Commissioner is hereby authorized to charge the above fee and any fee specifically authorized hereafter, or any missing or insufficient fee(s) filed, or asserted to be filed, or which should have been filed herewith or concerning any paper filed hereafter, and which may be required under Rules 16-18 (missing or insufficient fee only) now or hereafter relative to this application and the resulting Official document under Rule 20, or credit any overpayment, to our Deposit Account No. 21-0380, for which purpose a duplicate copy of this sheet is attached.

Submitted by:

Caroline Nash

Reg No 36,329

For: Charles H. Harris, Reg. No. 34,616
Attorney for Applicant
U.S. Army Medical Research and Materiel
Command, Attn: MCMR-JA
Fort Detrick, Maryland 21702-5012

Caroline M. Nash
Reg. No. 36, 329
Nash & Titus, LLC
3415 Brookeville Road
Suite 1000
Brookeville, Maryland 20833
Tel. No. (301) 924-9500
Fax. No. (301) 924-5883

NOTE: File in duplicate with post card receipt

CERTIFICATE OF MAILING BY "EXPRESS MAIL" (37 CFR 1.10)

Applicant(s): Reid, et al.

Docket No.

Army 103

~~Serial No.~~

U.S. Patent 5,693,343

Filing Date

5/16/94

issue date 12/2/97

Examiner

Craires, T.

Group Art Unit

1205

Invention: **MICROPARTICLE CARRIERS OF MAXIMAL UPTAKE CAPACITY BY BOTH M CELLS AND NON-M CELLS**

I hereby certify that this **Request for Filing Reissue Patent Application**

(Identify type of correspondence)


is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 in an envelope addressed to: The Assistant Commissioner for Patents, Washington, D.C. 20231

on **November 30, 1999**

(Date)

Caroline Nash

(Typed or Printed Name of Person Mailing Correspondence)



(Signature of Person Mailing Correspondence)

EJ843694054US

("Express Mail" Mailing Label Number)

Note: Each paper must have its own certificate of mailing.

IN RE THE
APPLICATION OF

ROBERT H. REID
JOHN E. VAN HAMONT
WILLIAM R. BROWN
EGAR C. BOEDEKER
CURT THIES

FOR REISSUE LETTERS PATENT OF
UNITED STATES PATENT NO. 5,693,343
FOR
“MICROPARTICLE CARRIERS OF MAXIMAL UPTAKE CAPACITY
BY BOTH M CELLS AND NON-M CELLS”

MICROPARTICLE CARRIERS OF MAXIMAL UPTAKE CAPACITY BY BOTH M CELLS AND NON-M CELLS

I CROSS REFERENCE

This application is a continuation-in-part of U.S. patent application Ser. No. 07/867,301 filed Apr. 10, 1992, now U.S. Pat. No. 5,417,986, which in turn is a continuation-in-part of U.S. patent application Ser. No. 07/805,721, filed Nov. 21, 1991, now abandoned, which in turn is a continuation-in-part of U.S. patent application Ser. No. 07/690,485 filed Apr. 24, 1991, now abandoned, which in turn is a continuation-in-part of U.S. patent application Ser. No. 07/521,945 filed May 11, 1990, now abandoned, which in turn is a continuation-in-part of U.S. patent application Ser. No. 07/493,597 filed Mar. 15, 1990, now abandoned, which in turn is a continuation-in-part of U.S. patent application Ser. No. 06/590,308 filed Mar. 16, 1984, pending.

II. GOVERNMENT INTEREST

The invention described herein may be manufactured, licensed and used by or for governmental purposes without the payment of any royalties to us thereon.

The file of this patent contains at least one drawing executed in color. Copies of this patent with color drawing(s) will be provided by the Patent and Trademark Office upon request and payment of the necessary fee.

III. FIELD OF THE INVENTION

The invention pertains in part to a method for preparing particle size distributions of microparticles of biodegradable polymers having the capacity to be maximally absorbed in both M cells and non-M cells in the Peyer's patches (PP) follicle-associated epithelium (FAE) and the villous epithelium region so that when the microparticles are used as carries of immunogens for oral immunization, the maximal conditions for uptake by gut lymphoid tissues will absorb any antigens so as to induce production of antibodies against diseases caused by the antigen or other enteropathogenic organisms, when using antigens encapsulated within biodegradable-biocompatible microspheres prepared by the process of the invention.

IV. BACKGROUND OF THE INVENTION

Infectious agents generally have their first contact with host organisms at a mucosal surface. Therefore, mucosal protective immune mechanisms are of key importance in preventing these agents from colonizing or penetrating the mucosal surface. It is apparent from past studies that a protective mucosal immune response can best be obtained by introduction of the antigen at the mucosal surface; however, parenteral immunization has not been an effective method to induce mucosal immunity. Antigen taken up by the gut-associated lymphoid tissue (GALT), primarily by the Peyer's patches stimulates T helper cells (T_H) to assist in IgA B cell responses or stimulates T suppressor cells (T_{KS}) to mediate the unresponsiveness of oral tolerance.

While particulate antigen appears to shift the responses towards the (T_H), soluble antigens favor a response by the (T_{KS}).

Although studies have demonstrated that oral immunization does induce an intestinal mucosal immune response, large doses of antigen are generally required to achieve sufficient local concentrations in the Peyer's patches. Further, unprotected protein antigens tend to be degraded or they complex with secretory IgA in the intestinal lumen.

One approach to overcoming the aforementioned problems is to homogeneously disperse the antigen of interest within the polymeric matrix of biodegradable, biocompatible microspheres that are specifically taken up by GALT. Eldridge, et al.¹ have used a murine model to show that orally-administered 1-10 micrometer microspheres consisting of polymerized lactide and glycolide, (the same materials used in resorbable sutures), were readily taken up into Peyer's patches, and that 1-5 micrometer sizes were rapidly phagocytized by macrophages. Microspheres that were 5-10 micrometers (microns) remained in the Peyer's patches for up to 35 days, whereas those less than 5 micrometer disseminated to the mesenteric lymph node (MLN) and spleen within migrating MAC-1⁺ cells.

¹Biodegradable Microspheres: Vaccine Delivery System For Oral Immunization, 1989, 146.

However, Eldridge, et al. used 50 μ m microspheres of poly (DL-lactide-co-glycolide) composed of molar parts of polymerized lactide and glycolide (85:15 DL-PLG), which biodegrades to completion in approximately 24 weeks after intramuscular injection.

Poly (DL-lactide-co-glycolide) composed of equal molar parts of polymerized lactide and glycolide (50:50 DL-PLG) is the more stable or least biodegradable, and biodegrades to completion after 25 weeks.

Therefore, there is a need extant in the biodegradable microsphere field to provide a method of producing poly (DL-lactide-co-glycolide) materials of 50:50 DL-PLG that is more biodegradable and capable of being taken up by both M cells and non-M cells in the Peyer's patches follicle-associated epithelium when used as microencapsulant as carriers for antigens for enteric immunization.

V. SUMMARY OF THE INVENTION

One object of the invention is to provide a method for producing microparticles of biodegradable-biocompatible microspheres having an average particle size distribution that maximizes uptake of the microspheres by both M cells and non-M cells, either in the villous epithelium or in the Peyer's patches follicle-associated epithelium so that, upon encapsulating antigens or other chemotherapeutic agents within these microspheres, large doses of antigen will not be required to achieve sufficient local concentrations in these regions of the intestines when these microparticles are used as carriers of immunogens for oral or other types of immunization.

A further object of the invention is to provide a method for producing microspheres composed of poly (DL-lactide-co-glycolide) having an average particle size distribution so as to maximize the uptake of these microspheres into the lymphoid tissue of the gut through uptake by both M cells and non-M cells, either in the villous epithelium or in the PP follicle-associated epithelium, in order to enable smaller doses of antigen to achieve sufficient local concentrations in these regions of the intestines when using the poly (DL-lactide-co-glycolide) as a carrier of immunogens for oral or other types of immunization.

A yet further object of the invention is to provide a method for producing an average distribution of particle sizes of the most stable or least biodegradable poly (DL-lactide-co-glycolide) having equal molar parts of polymerized lactide and glycolide (50:50 DL-PLG) so as to maximize uptake of microspheres of this copolymer by both M cells and non-M cells, either in the villous epithelium or in the PP follicle-associated epithelium when using this copolymer as a carrier of immunogens for oral or other types of immunization in mammals.

[illegible]

FIG. 1 shows that, during preparation of the microspheres, the spheres actually got larger as the emulsion time was increased.

FIG. 3 is a graph showing substitution of stable-viscosity machine oils or paraffin oils during the formation of the emulsion to accomplish sphere populations whose average sizes and volumes decreased with increasing emulsification times, in contrast to that which was observed for emulsions formed with paraffin oil as shown in FIG. 1.

FIG. 5 shows that reducing the viscosity of the paraffin oil by diluting it with heptane resulted in the formation of progressively larger spheres.

FIG. 6 shows that reducing the viscosity of the paraffin oil by diluting it with iso-octane resulted in the formation of progressively larger spheres.

FIG. 7 shows that when reducing the viscosity of the paraffin oil by diluting it with heptane using one second emulsification without an emulsion screen, resulted in the formation of progressively larger spheres.

FIG. 8 shows microsphere volume average versus emulsification time in paraffin oil.

FIG. 9 shows viscosity versus sphere diameter obtained with paraffin oil diluted with iso-octane.

FIG. 10 shows viscosity versus sphere diameter obtained with paraffin oil diluted with heptane.

FIG. 11 shows viscosity versus sphere diameter obtained with machine oils.

FIG. 12 is a color photograph of the flank region of the intestinal lymphoid follicle of a New Zealand white rabbit histochemically stained for acid phosphatase (red) and immunohistochemically stained for the MHCII antigen.

FIG. 13 is a color photograph of the flank region of the intestinal lymphoid follicle histochemically stained for alkaline phosphatase (red) and immunohistochemically stained for the MHCII antigen.

FIG. 14 is a color photograph of the flank region of the intestinal lymphoid follicle of a New Zealand white rabbit showing numerous microspheres of the poly (DL-lactide-

co-glycolide) composed of molar parts of polymerized lactide and glycolide (50:50 DL-PLG) in the company of MHCII-positive cells in lymphoid pockets in the Follicle Associated Epithelium (FAE), and wherein some of the
 5 microsphere particles are within the cells (arrows). In the lymphoid follicle, numerous MHUCII-positive cells are present, and some have microspheres associated with them (arrowheads).

FIG. 15 is a color photograph showing that both kinds of
 10 particles were taken up by the follicle-associated epithelium and entered the underlying lymphoid tissues of Peyer's patches (fluoresceinated microspheres are more easily visualized, and as a consequence they are shown in the photograph).

FIG. 16 is a color photograph showing the flank region of the previously illustrated intestinal lymphoid follicle and the adjacent villous stained for acid phosphatases (red) and CD43 (pan-T cell). Numerous CD43-positive cells are present in the FAE and in the lymphoid follicle. Microparticles in the FAE are in the company of CD43-positive cells in the lymphoid pockets, and some of the particles are within the cells (arrows). The CD43 cells are CD8- and CD4-negative, and Igu-positive cells are sparse in the FAE. The microparticles have not entered the epithelium of the villous
 25 (v) adjacent to the lymphoid follicle, although some are present nearby in the lumen.

FIG. 17 is an immunofluorescence micrograph of the previously illustrated lymphoid follicle. The fluorescein-labeled microspheres are present mostly in the flank region
 30 of the FAE (lower area of the photograph), with declining numbers present in the more apically located regions.

FIG. 18 is a color photograph showing the lymphoid follicle of a Peyer's patch of a New Zealand white rabbit stained for vimentin. The polymerized lactide and glycolide particles appear principally in the FAE area and are practically non-existent in the villous area.

FIG. 19 is a color photograph of the lymphoid follicles of the New Zealand white rabbit's intestines showing the pan-T cell markup stained for CD43. The view shows the villous
 40 epithelium, the lamina propria, the location of the copolymer (PLGA) particles and the CD43-positive cells.

VII. DETAILED DESCRIPTION OF THE INVENTION

45 Use of the emulsion viscosity as the means for controlling the average particle size distribution of polymerized lactide and glycolide microspheres has utility in manufacturing oral and injectable vaccines as well as for use in devices for
 50 sustained drug and antibiotic delivery. Preparation of the microspheres was accomplished by a modification of the solvent extraction process to control the sphere size by altering the viscosity of the emulsion either by: 1) predilution of the emulsion oil with an extractant solvent; 2)
 55 adding thickening agents such as polybutylene to the emulsion oil to deliberately increase its viscosity; 3) through use of oils with predefined viscosities for preparation of the emulsion; or 4) through deliberately adjusting the viscosity of the paraffin oil by preheating it to a temperature which
 60 yields the desired viscosity, taking care that the emulsion time is kept sufficiently short so as to prevent a significant temperature increase during the emulsification process.

It has been found that the oil viscosity is the primary process parameter for controlling the sphere diameter, and
 65 that variation in screen and rotor dimensions, emulsification speed and time only exhibit negligible effects on the outcome of the diameter of the microspheres.

Solvent Extraction

Materials

80 mg sucrose

20 mg purified antigen/active

Preparation of the Antigen-Sucrose Matrix

The dispersion is then flash frozen by gently swirling the vial (without the cap) in liquid nitrogen for about one half of an hour.

After about 1000 minutes the temperature is elevated to about +5 degrees for 500 minutes (8.33 hours) and then elevated to about +20 degrees for 1000 minutes (16.67 hours), and the vial is removed.

The PLG is removed from the freezer and allowed to come to room temperature.

About 2.8 g of acetonitrile is weighed into a 20 ml capacity glass vial and set aside.

After the polymer reaches room temperature, about 1.0 g of the polymer is added to the vial of acetonitrile and a sonicator bath until all of the polymer has dissolved (5-10 minutes).

Preparing the Homogenizer

3.2 g of acetonitrile is weighed in a plastic vial for washings during homogenization.

1.5 g of acetonitrile is weighed into another vial and added to the earlier prepared freeze-dried sucrose-antigen matrix and mixed until it becomes a milky white slurry. The slurry is homogenized at maximum speed for one minute and the 3.2 g of acetonitrile is used to wash the sides of the vial and homogenizer tip, after which the slurry is again homogenized for one minute at maximum speed.

The mixture is separated into two parts by weight by weighing 2.4 g into another 20 ml plastic vial.

The polymer solution prepared earlier is added to one of the vials of homogenized sucrose-antigen and the vial is placed in a sonicator bath for about 2 minutes to ensure proper mixing.

The homogenizer is set up with the rotor and fine emulsion screen and the following materials are weighed out: 400 g of light mineral oil in a 600 ml glass beaker and 2500 g of heptane in a 4000 ml propylene beaker. A beaker of heptane is placed under the mixer and a propeller is placed about two-thirds of the way down into the heptane, after which the mixer is started at about 450 rpm.

A Masterflex mode #7550-60 peristaltic pump with pump head model #7518-00 with PharMed tubing, size 16 is set up. The pump speed is set at 300 ml/min after which one end of the tubing is placed into the beaker of heptane.

Approximately 175 ml of the mineral oil is poured into the jacket beaker and the homogenizer head is dipped at 15 degrees into the oil to coat it, after which water is circulated through the beaker.

- 5 The polymer/sucrose-antigen solution is poured into the beaker and the vial is rinsed with about 5 ml of mineral oil, and the rinse is added to the beaker.

The homogenizer head is placed into the liquid and then turned to its maximum setting for 3 minutes.

- 10 At the end of three minutes the other end of the tubing is placed into the jacketed beaker and the peristaltic pump is started. When the liquid level has dropped to the level of the homogenizer head, the homogenizer is turned off and pumping is continued until all of the liquid has been pumped to the
15 heptane after which the heptane is left stirring for 30 minutes.

- Using fresh tubing, the heptane is pumped into centrifuge bottles and centrifuged for 5 minutes at 3000 rpm. 20 degrees celsius. The supernatant is pumped into waste
20 bottles and the sediment is rinsed with heptane (it may be necessary to sonicate the sample for 1 to 5 minutes to break up the sediment).

- The supernatant is pumped into the waste bottle and washed with fresh heptane until all the microspheres are in
25 the one tared 50 ml centrifuge tube. This tube is then centrifuged for 5 minutes and washed with fresh heptane three times.

- After the final wash and centrifuge cycle, the supernatant is pumped into the waste bottle and the microspheres are air
30 dried with a slow air current for about 5 minutes, and the tube is placed in the vacuum oven at room temperature and left overnight.

- The microspheres are removed from the vacuum oven and weighed, after which about 1 mg of the microspheres is put
35 in a 1.5 ml centrifuge tube for evaluation.

Evaluation of Microspheres

- About 1 ml of 1% Tween 80 in water is added to the 1 mg of microspheres in the 1.5 ml centrifuge tube, and the tube
40 is sonicated for about 1 minute.

- One drop of the dispersion is placed on a glass slide and a coverslip is placed over it. The slide is then placed under a calibrated optical microscope and examined under 100× magnification using a standard oil immersion technique.
45 Using the precalibrated eyepiece micrometer, the diameter of 150 randomly chosen microspheres is determined. (Under 100× magnification, 1 division on the micrometer is equal to 1 micron.)

- The numbers are then entered into a Lotus spread sheet program to determine the average size distribution of the
50 particles.

- The prior art extraction procedure for production of poly-lactide: glycolide microencapsulated oral vaccines is based on disbursal of a highly concentrated solution of
55 polymer and acetonitrile into oil, followed by extraction of the acetonitrile and oil with heptane.

- The procedure of the invention requires high energy shear to disburse the viscose polymer solution. This high shear process resulted in the generation of major heat change
60 which caused the mineral oil's viscosity to change significantly.

- As a result, small increases in shear time or minor differences in the emulsifier's rotor dimensions which increased shear, resulted in increased microsphere
65 diameters, as can be seen from the graph of FIG. 16.

The substitution of stable viscosity machine oils for paraffin oil during the formation of the emulsion resulted in

sphere populations whose average sizes and volumes decreased with increasing emulsification times.

This result can be seen in Table I, which is in contrast to the data showing microsphere volume average versus emulsification time and paraffin oil (FIG. 16).

TABLE I

C/S*	Emulsification Time			
	0.75 minutes		3.0 minutes	
	V.A.**	D.A.***	V.A.	D.A.
36	2.8	0.9	2.4	0.9
50	7.3	2.9	6.8	3.2
65	7.9	3.3	6.9	2.9
72	4.9	1.0	3.0	0.9
80	2.4	1.1	1.4	0.9

* = Centistokes

** = Volume Average

*** = Diameter Average

Both paraffin emulsions and machine oil emulsions underwent similar temperature increases during the emulsion process, and the differences between these two oils appears to be due to maintenance of a relatively constant emulsion viscosity by the machine oils. At a constant viscosity, increased homogenization time appears to have resulted in a progressively finer dispersal of the polymer-acetonitrile solution into the oil. Viscosity breakdown in the paraffin oil appears to have allowed particles to recombine as the emulsion temperature increased.

Reducing the viscosity of the paraffin oil by diluting it with either heptane or iso-octane resulted in the formation of a progressively larger spheres as can be seen in Table 2.

TABLE 2

Sphere Diameters Resulting From Dilution of the Emulsion Oil			
Solvent/ Oil Mixture	Diameter Averages in U		
	H*	IO**	IO***
1/2	11.0	4.6	9.7
1/4	2.6	5.0	3.3
1/8	1.5	2.2	2.6
1/16	1.2	1.4	0.6
No Solvent	1.0	1.0	0.6

* = Heptane Diluent.

** = Iso-octane.

*** = 2nd Series of Iso-octane Batches Employing Reduced Shear Forces

The results of these tables show that sphere size can be controlled by altering the viscosity of the emulsion oil through its pre-dilution with an extractant solvent, provided that the emulsion time is kept sufficiently short so as to prevent a significant temperature increase during the emulsification process.

The data in Table 1 shows a relationship between the microsphere size and oil viscosity in that, microsphere size increased as oil viscosity increased from 36 to 65 centistokes and then decreased from 65 to 80 centistokes, which appears to indicate a bell-shaped sphere size distribution as viscosity increased.

FIG. 17 shows viscosity versus sphere diameter obtained with paraffin oil diluted with iso-octane.

FIG. 18 shows viscosity versus sphere diameter obtained with paraffin oil diluted with heptane.

FIG. 19 shows viscosity versus sphere diameter obtained with machine oils.

A histochemical and immunohistochemical analysis of the uptake of PLG and polystyrene microparticles by Peyer's patches from a New Zealand white rabbit was conducted using the poly (DL-lactide-co-glycolide) copolymer in which the molar parts of polymerized lactide and glycolide were 50:50, as prepared according to the modified solvent extraction process of the invention.

- 10 Fluorescent polystyrene microspheres were also used as a comparison to test these microparticles as carriers of immunogens for oral immunization, and to ascertain or determine the actual location of their uptake by gut lymphoid tissues, and to ascertain which tissues were engaged in the uptake.

The study also served in part to ascertain if encapsulation may protect the antigens from proteolytic degradation in the gut lumen and facilitate their uptake and retention in the intestinal lymphoid tissues, as a thorough understanding of the fate of ingested antigen-containing microparticles is important in using antigens which have been microencapsulated for enteric immunization strategies.

VIII. METHOD

- 30 Fluorescent polystyrene microspheres and unlabelled poly (lactide-co-glycolide) microspheres of diameters of 0.5, 1, and 2 μ m were instilled into the lumens of in situ rabbit intestinal loops.

After a period of between about 1 to 2 hours, the loops were removed, and sections were cut and reacted histochemically for acid (AcP), phosphatase and immunohistochemically in a biotin-streptavidin method with several monoclonal antibodies to the rabbit lymphoid cell antigens.

The rabbits were anesthetized New Zealand white rabbits and the dimensions of the intestinal loops were 2 cm (containing Peyer's patches) and the tissue blocks were excised and fixed in periodate-lysine-paraformaldehyde.

- 50 The results of these tests show that both kinds of particles (0.5-2 μ m) were taken up by the Peyer's patches.

However, the particles of copolymer from the invention process principally taken up in the Peyer's patch region have a volume average of about 1.0 to about 7.0 micrometers as the particle size distribution.

- 60 Particles of copolymer from the invention process principally taken up in the villous epithelium of the intestines have a volume average of from about 0.5 to about 2.0 micrometers.

Tables 3 and 4 show respectively, the particles used when testing placement in Peyer's patch and villous regions.

Microspheres: Particle Size Distribution by Microscopy

Stage magnification: 100 ×									
Calibration: 1 div = (u)									
1.00									
between or equal to 5-10 u									
% volume % number									
73.25 13.53									
greater than 10 u									
% volume % number									
0.00 0.00									
Reading	Frequency Dia.,	u	d ³	FD ³	% Vol. Dist.	.01*F*d	F*D	% Num Dist.	
1	40	1	1	40	0	0	40	27	between or equal to 1-5 u
2	41	2	8	329	3	0	82	27	% volume % number
3	25	3	27	675	6	2	75	17	26.75 86.67
4	13	4	64	832	7	4	52	9	
5	11	5	125	1375	11	14	55	7	
6	9	6	216	1944	16	35	54	6	
7	5	7	343	1715	14	48	35	3	
8	1	8	512	512	4	22	8	1	
9	1	9	729	729	6	44	9	1	
10	4	10	1000	4000	33	329	40	3	
Total	150	55	3025	12150	85	498	450	100	
average		5.50	6.71		2.15		3	2.15	
S.D.		2.87	6.88		2.08		2.85	2.12	

AGGREGATION DATA		
Particle Size Distribution	Vol % Aggregated	0.00 Weight %
Volume Average (u)	Number % Aggregated	0.00 number %
Number Average (u)	Avg Part Size Aggregated	ERR
	Avg # of Part per Aggregate	ERR

Microspheres: Particle Size Distribution by Microscopy

[illegible]

The significance of what particle size distribution of the copolymer prepared according to the invention process is taken up in the villous epithelium section of the intestine is that, for oral administration of a vaccine (especially when no booster vaccine is administered), the antigen must principally be uptaken by the villous epithelium region, which is more than 90% of the area of the intestine needed for effective immunization. On the other hand, the fact that

The following information obtains from a immunohistochemistry basis:

IMMUNOHISTOCHEMISTRY

Antibodies

IMMUNOHISTOCHEMISTRY		
Antibodies: Monoclonal Antibody	Source	Antigen Recognized
V9	Biomed	Vimentin (M cell marker)
L11-35	Serotec	CD43 (pan T cell)
45-3	Spring Valley	MHC II
Ken-4	Spring Valley	CD4
12C7	Spring Valley	CD8
NRBM	Serotec	Ig u chain
Procedure: Biotin-streptavidin method		

Uptake was greatest along the flanks of the follicles, where M cells (demonstrated by anti-vimentin MAb) were most numerous. While the particles were sometimes present within M cell cytoplasm, they were much more numerous in the lymphocyte pockets of the M cells.

In the pockets, the particles were intermingled with cells that were CD43+, CD8-, CD4-, Igu-, and MHC II+.

The results showed that, occasionally, the particles were present within large AcP+ cells in the pockets. In the follicular tissue beneath the M cell-rich epithelium, particles were very numerous in the vicinity of MHC II+ cells and occasionally within the large AcP+ cells.

Unexpectedly, microspheres also entered non-M cell epithelium cells, especially in the domes. These cells were vimentin negative AcP+. Microparticles were sparse or absent in the subepithelial tissue beneath the cells.

As a result of these tests, it became clear that both M cells and non-M cells in the rabbit PP follicle-associated epithelium can take up certain microparticles. Only the M cells may be capable of permitting migration of the particles to adjacent cells.

Microparticles taken up by the M cells appear to migrate to lymphocyte pockets richly populated with MHC II+ cells and CD8-/CD4- T cells, as well as to a certain extent to AcP+ phagocytic cells.

What is claimed is:

1. In a solvent extraction process for preparing microspheres of an antigen containing biodegradable poly(DL-lactide-co-glycolide), the improvement comprising:
 - 5 preparing a lyophilized antigen-sucrose matrix; adding acetonitrile solvent to the antigen-sucrose matrix to form a solution;
 - 10 preparing a solution of a biodegradable poly (DL-lactide-co-glycolide) polymer by adding acetonitrile solvent to the polymer;
 - 15 adding the biodegradable poly (DL-lactide-co-glycolide) polymer acetonitrile solution to the antigen-sucrose acetonitrile solution;
 - 20 adding an oil to the poly (DL-lactide-co-glycolide) polymer-sucrose-antigen solution to form an emulsion having a controlled viscosity, that corresponds to a predetermined average particle size of distributions of microspheres of poly (DL-lactide-co-glycolide) biodegradable polymers of from about 0.5 to about 7.0 micrometers;
 - 25 centrifuging the emulsion of controlled viscosity and removing a supernatant to obtain microspheres of the predetermined range of particle size distributions.
2. The process of claim 1, wherein the oil is selected with a predefined viscosity to form the microspheres.
3. The process of claim 1, wherein a thickening agent is
30 added to the oil to increase its viscosity.
4. The process of claim 1, wherein the oil is prediluted with an extractant solvent.
5. The process of claim 1, wherein the oil is a paraffin oil in which the viscosity is adjusted by preheating to a tem-
35 perature of desired viscosity.
6. The process of claim 1, wherein relative ratios between the lactide and glycolide is 50:50.
7. The process of claim 1, wherein the average particle
40 size distribution is from about 1.0 to about 2.0 micrometers.

* * * * *

1660667-1
/ Claim 19. A composition comprising an encapsulating-microsphere of a biodegradable polymer having an average particle size distribution wherein a majority of the microspheres will be taken up by a villous epithelium section of an intestines of a mammalian subject when administered.

Claim 20. The composition of claim 19, wherein said biodegradable polymer comprises a poly (DL-lactide-co-glycolide) copolymer.

Claim 21. The composition of claim 19, wherein said average particle size distribution is about 0.5 to about 2.0 micrometers.

1660667-1
/ Claim 22. A composition comprising an encapsulating-microsphere of a biodegradable polymer having an average particle size distribution wherein a majority of the microspheres will be taken up by a Peyer's patch section of an intestines of a mammalian subject when administered.

Claim 23. The composition of claim 22, wherein said biodegradable polymer comprises a poly (DL-lactide-co-glycolide) copolymer.

Claim 24. The composition of claim 22, wherein said average particle size distribution is about 1.0 to about 7.0 micrometers.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION OF

REID et al. REISSUE of U. S. Patent No. 5,693,343

Appln. No.: TBA

Group Art Unit: TBA

Filed: Herewith

Examiner: TBA

Title: MICROPARTICLE CARRIERS OF MAXIMAL UPTAKE CAPACITY BY
BOTH M CELLS AND NON-M CELLS

LETTER REQUESTING TRANSFER OF THE DRAWINGS
FROM THE PATENT FILE UNDER MPEP 1413

Hon. Commissioner of Patents
and Trademarks
Washington, D.C. 20231

Sir:

Please transfer the drawings upon which the original patent was issued to this reissue application in lieu of new drawings. No changes whatsoever are to be made in the drawings. Temporary drawings which consists of a copy of the printed drawings of the patent or photoprint of the original drawings of the size required for original drawings are submitted with this reissue application.

Respectfully submitted,

Caroline Nash Reg. No. 36,329

Date: *Nov. 30, 1999*

For: Charles H. Harris, Reg. No. 34,616
Attorney for Applicant
U.S. Army Medical Research and Materiel
Command, Attn: MCMR-JA
Fort Detrick, Maryland 21702-5012

Caroline M. Nash, Reg. No. 36, 329
Nash & Titus, LLC
3415 Brookeville Road
Suite 1000
Brookeville, Maryland 20833
Tel. No. (301) 924-9500
Fax. No. (301) 924-5883

FIG. 1




FIG. 2





FIG. 3




EMULSION OIL VISCOSITY IN CENTISTOKES @ 40°C

X—X - 0.75" EMULSIFICATION TIME

O---O - 3.0" EMULSIFICATION TIME

FIG. 4



* - VISCOSITY ON OIL IN CENTISTOKES @ 40°C

FIG. 5




FIG. 6





FIG. 7



ISO-OCTANE / ISO-OCTANE + OIL RATIO *

*ONE SECOND EMULSIFICATION WITHOUT AN
EMULSION SCREEN

FIG. 8





FIG. 9






FIG. 11





FIG 12




FIG 13



660611-125460

FIG 14




660E.T. T28F5460

FIG. 15



660577-1-23-5460

FIG 16



560577-125460

FIG. 17






FIG. 17

FIG. 18



650EFT" T2E T5460

FIG. 19



650E.T. T28T5460

DECLARATION AND POWER OF ATTORNEY FOR REISSUE APPLICATION

We, Robert H. Reid, John E. van Hamont, William R. Brown, Egar C. Boedeker, and Curt Thies, hereby declare that we are citizens of the United States and that our residences are as stated below next to our names. We believe that we are the original joint inventors of the invention entitled MICROPARTICLE CARRIERS OF MAXIMAL UPTAKE CAPACITY BY BOTH M CELLS AND NON-M CELLS described and claimed in our original application No. 08/242,960, filed May 16, 1994, and the resulting United States Patent No. 5,693,343 which issued December 2, 1997 ("the '343 patent"), for which priority was claimed based on Ser. No. 867,301, filed April 10, 1992, Pat. No. 5,417,986 which is a continuation in part of Ser. No. 805,721, filed Nov. 21, 1991, abandoned, which is a continuation in part of Ser. No. 690,485, filed April 24, 1991, abandoned, which is a continuation in part of Ser. No. 521,945, filed May 11, 1990, abandoned, which is a continuation in part of Ser. No. 493,597, filed March 15, 1990, abandoned, which is a continuation in part of Ser. No. 590,308, filed March 16, 1984, and for which invention a reissue patent is solicited.

We do not know and do not believe that the invention of the '343 patent was ever known or used in the United States before our invention thereof. Furthermore, we do not know and do not believe that the invention was patented or described in any publication in any country before our invention thereof, or more than one year prior to the original application. We do not know and do not believe that the invention was in public use or on sale in the United States more than one year prior to the original application. To the best of our knowledge and belief, this invention has

not been patented or made the subject of an inventors' certificate in any country foreign to the United States prior to the date of the original application on an application filed by us or our legal representatives or assigns more than 12 months before our original application.

We have reviewed and understand the contents of the attached specification, including the claims, as amended by the addition of new claims 8-24. We acknowledge the duty to disclose information of which we are aware and which is material to the examination of the application in accordance with 37 C.F.R. §§ 1.56(a) and 1.175(a) (7).

We believe that through error, without any deceptive intent, the '343 patent is partially inoperative or invalid by reason of the patentee claiming less than the patentee had the right to claim in the patent. In particular, there is a possible defect in that the patentees claimed less than patentees had the right to claim in the patent.

This possible error arose without deceptive intent during prosecution of the application before the United States Patent and Trademark Office. Upon reviewing the issued claims we realized that the claims did not cover all of the subject matter that the we believe we are entitled to.

Therefore, by reason of the above-described error, Applicants believe the original patent to be partly inoperative or invalid by reason of the patentees claiming less than patentees had the right to claim in the patent and the claims are possibly not broad enough to cover all aspects of the invention disclosed in the patent. The possible invalidity of the patent resulted from a failure by ourselves, the assignee and counsel to realize the totality of the subject matter that should have been claimed. By this reissue application, the identified errors are believed to be corrected.

Reid, et al. Reissue Application of U.S. Patent 5,693,343**Inventor's Name:** John E. van Hamont**Inventor's Residence:** 239B Barnard Loop, West Point, New York 10096, USA**Inventor's Citizenship:** U.S.A.**Inventor's Signature:** John E. van Hamont**Date Signed:** 10 November 1999

Inventor's Name: Edgar C. Boedeker

Inventor's Residence: 552 Maynadier Lane, Crownsville, MD 21032, USA

Inventor's Citizenship: U.S.A.

Inventor's Signature: _____

Date Signed: 10/21/99

CONFIDENTIAL

All errors which are being corrected in the present reissue application up to the time of filing of this declaration arose without any deceptive intention on the part of the applicants.

Wherefore we request that we may be allowed to surrender, and we hereby offer to surrender, said U.S. Letters Patent No. 5,693,343 and request that Letters Patent be reissued to ourselves and the assignee, The United States of America as represented by the Secretary of the Army, for the same invention upon the foregoing amended reissue application.

We hereby declare further that all statements made herein of our knowledge are true and that all statements made on information and belief are believed to be true. We further declare that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under § 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

We hereby appoint Charles H. Harris (Reg. No. 34,616) of the U.S. Army Medical Research and Materiel Command, 504 Scott Street, ATTN: MCMR-JA, Fort Detrick, MD 22702 and Caroline Nash (Reg. No. 36,329) and Marlana K. Titus (Reg. No. 35,843) of Nash & Titus, LLC, 3415 Brookeville Road, Suite 1000, Brookeville, Maryland 20833, (301) 924-9500 or (301) 924-9600 (all communications are to be directed to Nash & Titus, LLC) individually and collectively as our attorneys to prosecute this application and to transact all business in the Patent and Trademark office connected therewith and with the resultant Patent.